

Can extensions of the standard model inform a low-energy theory of quantum gravity?

Robert N. C. Pfeifer*

School of Mathematics and Physics, The University of Queensland, Brisbane, Queensland 4072, Australia

(Dated: June 4, 2009)

The search for a quantum theory of gravity has largely focused on *de novo* approaches such as string theory and loop quantum gravity, with the assumption that unification with other forces of the standard model is only relevant at very high energy scales. This paper asks instead whether we can learn anything about a quantum theory of gravity by extending the standard model, attempting in some limited regime to reproduce appropriately gravity-like phenomena. As a concrete example the gauge group of the electroweak interaction is extended to $GL(3, \mathbb{R})$, accompanied by modification of the Higgs interaction. Although containing extra bosons which may be unphysical, the toy model presented here is notable in that gauging of the space-time symmetry group permits derivation of the Schwarzschild metric in the vicinity of an irrotational, spherically symmetric body, along with an explicit expression for the value of G . Evaluation of the expression gives $G = 6.675\,514\,4(15) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$, in agreement with the NIST/CODATA recommended value of $G = 6.674\,28(67) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$ to within two standard deviations of the experimental result. These results strongly indicate that this model, and perhaps others like it, may indeed be able to inform our construction of an effective low-energy theory of quantum gravity.

PACS numbers: 04.60.-m, 12.60.Cn, 12.10.-g, 04.50.-h

I. INTRODUCTION

The search for a quantum theory of gravity is one of the greatest challenges facing theoretical physics today, requiring mathematics far beyond the standard quantum field theory which suffices to describe the other four physical forces. The approaches of both string theory [1, 2] and loop quantum gravity [3, 4] attack this problem head-on, aiming to construct a quantum theory of gravity in isolation, and leaving unification with the rest of the standard model as a problem for the future.

Nevertheless, it is possible there is some benefit in asking, even at this early stage, how a quantum theory of gravity will relate to the rest of the standard model. One way to pursue this goal might be the construction of extended portions of the standard model which reproduce in some limited domain behaviours which we might associate with gravitation. From such extended theories we might gain insight into the low energy behaviour of a full quantum theory of gravity.

In this paper, I present a modification to the electroweak and Higgs sectors of the standard model, introducing several extra bosons which expand the internal symmetry group to either $SU(3)$ or $GL(3, \mathbb{R})$. The extra bosons have never been detected, and no attempt is made to claim that they necessarily ever will be (though prospects for detecting them if they are real are discussed in Sections IV A and V D). Of greater interest are the consequences of choosing a gauge on the combined internal and external symmetry groups $GL(3, \mathbb{R}) \otimes P_{3,1}$, which may be done in such a manner as to eliminate some of the extra bosons which have been introduced. Moreover,

we find that the metric of this co-ordinate frame in the vicinity of an irrotational, spherically symmetric object is the Schwarzschild metric, and that in this model we may calculate the value of Newton's gravitational constant, G . The result is $G = 6.675\,514\,4(15) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$, which lies within 1.8σ of the NIST/CODATA recommended value of $6.674\,28(67) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$ [5, 6]. Consequently, in the vicinity of an irrotational, spherically symmetric massive body the use of this prescription for gauging local Lorentz invariance is consistent with Einstein's general theory of relativity.

Whether this model is physical, or is merely a convenient approximation to some more fundamental theory of gravity with the extra bosons acting as proxy for a more sophisticated underlying interaction, the precision with which it calculates the value of G suggests that when the true quantum theory of gravity is found, its behaviour in the low-energy limit will in some way resemble the model presented in this paper. Which of the physical hypotheses employed in this model will turn out to be true, and which are merely convenient fictions, only time will tell.

II. THE MODEL

A. Lagrangian

Consider the following Lagrangian density:

$$\mathcal{L} = \bar{\Phi} \not{D} \Phi - \frac{1}{4} \text{Tr} F^{\mu\nu} F_{\mu\nu} \quad (1)$$

$$\Phi = \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix} \quad (2)$$

$$\mathcal{D}_\mu = \mathbb{I}_3 \otimes \partial_\mu - i f b_\mu^a C_a \quad (3)$$

$$F_{\mu\nu} = \mathcal{D}_\mu b_\nu^a C_a - \mathcal{D}_\nu b_\mu^a C_a. \quad (4)$$

*pfeifer@physics.uq.edu.au

In this expression, ϕ_1 , ϕ_2 , and ϕ_3 are left-handed anticommuting Weyl spinor fields, representing fermionic particles. Denoting the space of left-handed Weyl spinors $D(\frac{1}{2}, 0)$, we may represent the state of these three fields by the vector Φ on $D(\frac{1}{2}, 0)^{\otimes 3}$. The Lagrangian density (1) is invariant under the global action of the gauge group $SU(3)$ acting on Φ , and the Euler–Lagrange equations of motion are invariant under the action of $GL(3, \mathbb{R})$. Promoting the symmetry from global to local, matrices C_a represent a basis of the corresponding Lie algebra, either $su(3)$ or $gl(3, \mathbb{R})$ respectively, $\mathcal{D}_\mu$ defines the covariant derivative, and the b_μ^a correspond to the resulting vector bosons, coupling with interaction strength f . Their propagator is introduced in the usual manner, in terms of a field strength $F^{\mu\nu}$. We will adopt units such that $\hbar = c = 1$, and suppress spinor indices where this may be done without ambiguity. Mass takes units of inverse length. The metric signature is $(-1, 1, 1, 1)$ and the sigma matrices are

$$\bar{\sigma}^\mu = \{\mathbb{I}_2, \vec{\sigma}\} \quad (5)$$

$$\sigma^\mu = \{\mathbb{I}_2, -\vec{\sigma}\}. \quad (6)$$

Let us also postulate the existence of a spontaneous symmetry breaking mechanism which, although not explicitly specified, has the effect of attributing a ground-state value to $\bar{\Phi}(\bar{\sigma}^\mu b_\mu^a C_a)\Phi$ which is non-zero and homogeneous in space and time. We assume that this hypothetical symmetry breaking process does not favour any particular orientation with respect to either external or internal symmetry groups.

We now employ our local $SU(3)$ symmetry to perform a choice of gauge, first mixing ϕ_1 , ϕ_2 , and ϕ_3 such that

$$\bar{\Phi}(\mathbb{I}_3 \otimes \bar{\sigma}^\mu)\Phi = \bar{\phi}_3 \bar{\sigma}^\mu \phi_3. \quad (7)$$

From this it follows that $\bar{\phi}_1 \bar{\sigma}^\mu \phi_1 = -\bar{\phi}_2 \bar{\sigma}^\mu \phi_2$, and this is satisfied by ϕ_1 and ϕ_2 of the form

$$\phi_1 = q e^{i\theta} \phi_2, \quad 0 \leq \theta < 2\pi, \quad q \in \Lambda, \quad q^* q = -1, \quad (8)$$

where Λ denotes the space of Grassman numbers. Consequently, we may perform a further $SU(2)$ mixing of ϕ_1 and ϕ_2 such that

$$\bar{\phi}_1 \bar{\sigma}^\mu \phi_3 = \bar{\phi}_2 \bar{\sigma}^\mu \phi_3 = 0. \quad (9)$$

The requirement that the fermion and boson fields be in equilibrium in the ground state then ensures that the boson fields to which $\bar{\phi}_1 \bar{\sigma}^\mu \phi_3$, $\bar{\phi}_2 \bar{\sigma}^\mu \phi_3$, $\bar{\phi}_3 \bar{\sigma}^\mu \phi_1$, and $\bar{\phi}_3 \bar{\sigma}^\mu \phi_2$ couple are also zero.

In the ground state, all fields will satisfy equations of motion which are homogeneous in space and time. Where coupling to non-zero vacuum expectation values occurs, these equations of motion will be those of a massive particle, and minimisation of the Lagrangian favours a solution to these equations of motion where the rest frame is uniform throughout space–time. However, we will assume that the universe is in a low energy state slightly

above the ground state (though still within the broken symmetry regime), and consequently we will permit this rest frame to vary smoothly in space and time.

Finally we must extend our choices of gauge to states other than the ground state, introducing some notion of particle excitations on the vacuum. Let us therefore admit regions where the field strengths vary from the vacuum expectation value:

$$\bar{\Phi}(\mathbb{I}_3 \otimes \bar{\sigma}^\mu)\Phi = [\bar{\Phi}(\mathbb{I}_3 \otimes \bar{\sigma}^\mu)\Phi]_{\text{bg}} + [\bar{\Phi}(\mathbb{I}_3 \otimes \bar{\sigma}^\mu)\Phi]_{\text{fg}} \quad (10)$$

$$b_\mu^a C_a = [b_\mu^a C_a]_{\text{bg}} + [b_\mu^a C_a]_{\text{fg}} \quad (11)$$

We extend our choice of gauge by dividing the fields into background and foreground components in the manner of (10), where $[\]_{\text{bg}}$ represents the local background field attributable to the non-zero vacuum expectation value, $[\]_{\text{fg}}$ represents the deviation from this value, and we choose our orientation of ϕ_3 so as to minimise the value of $[\bar{\phi}_3 \bar{\sigma}^\mu \phi_3]_{\text{fg}}$ at every point in space and time. The deviations from the vacuum state represented by $[\bar{\Phi}(\mathbb{I}_3 \otimes \bar{\sigma}^\mu)\Phi]_{\text{fg}}$ and $[b_\mu^a C_a]_{\text{fg}}$ represent our notion of particle or anti-particle excitations, depending on their indices and sign.

The generalisation of our second gauge condition is chosen to enable us to eliminate some of the boson fields in Sec. VB. Where the boson fields coupling to $\bar{\phi}_1 \bar{\sigma}^\mu \phi_3$ and $\bar{\phi}_3 \bar{\sigma}^\mu \phi_1$ are non-zero, we perform an $SU(2)$ mixing on ϕ_1 and ϕ_2 such that after mixing,

$$\bar{\phi}_1 \bar{\sigma}^\mu \phi_3 = \bar{\phi}_3 \bar{\sigma}^\mu \phi_1. \quad (12)$$

In the ground state, this is consistent with condition (9).

B. Particle identities

To relate this theory to the electroweak sector of the standard model, we must first choose a basis for our internal symmetry group. A convenient choice will be

$$\begin{aligned} C_1 &= \frac{2}{\sqrt{6}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & 0 \\ 0 & 0 & -\frac{1}{2} \end{pmatrix} & C_2 &= \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ C_3 &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} & C_4 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ C_5 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} & C_9 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ C_6 &= C_2^\dagger & C_7 &= C_3^\dagger & C_8 &= C_5^\dagger, \end{aligned} \quad (13)$$

if the symmetry group is $GL(3, \mathbb{R})$, or with C_9 omitted if the symmetry group is $SU(3)$. We now identify

$$\begin{aligned} g &= \sqrt{2} f & \phi_2 &= e_L & \phi_3 &= \nu_e \\ b_\mu^1 &= X_\mu & b_\mu^4 &= W_\mu^3 & b_\mu^5 &= W_\mu \\ b_\mu^2 &= H_\mu & b_\mu^3 &= G_\mu & b_\mu^9 &= N_\mu. \end{aligned} \quad (14)$$

where e_L is the left-handed electron, ν_e the left-handed electron neutrino, and capital letters represent gauge bosons. The W boson is familiar, and the W^3 and X bosons will be mixed to yield the photon and Z boson in the usual manner. We shall see shortly that the H boson plays a Higgs-like role in giving mass to the electron, the G boson is indirectly involved in the gravitational interaction, and the N boson (if it exists) is a weakly interacting massive neutral boson, coupling only to the gravitational field, W and $W^\dagger$ bosons, and the neutrino, making it a possible candidate for WIMP dark matter. The apparent additional weak interaction involving the G and $G^\dagger$ bosons will be eliminated in Sec. VB.

We have yet to assign an identity to left-handed spinor ϕ_1 . Instead of doing so directly, we shall make use of an isomorphism

$$F : \bar{\lambda} \rightarrow \rho \quad (15)$$

between the space of charge conjugate left-handed spinors $\overline{D(\frac{1}{2}, 0)}$ and the space of right-handed spinors, $D(0, \frac{1}{2})$. For the explicit representation

$$\lambda = \begin{pmatrix} \lambda_1 + i\lambda_2 \\ \lambda_3 + i\lambda_4 \end{pmatrix} \quad \lambda_i \in \Lambda, \quad (16)$$

this is given (up to a freedom of sign on λ_2 and λ_4) by

$$F : (\lambda_1 - i\lambda_2 \quad \lambda_3 - i\lambda_4) \rightarrow \begin{pmatrix} \lambda_1 - i\lambda_2 \\ \lambda_3 - i\lambda_4 \end{pmatrix}. \quad (17)$$

We now identify the charge conjugate $\bar{\phi}_1$ with the right-handed electron e_R ,

$$\bar{\phi}_1 = e_R, \quad (18)$$

and in light of this isomorphism, we may use e_R to denote either a right-handed spinor or the charge conjugate of a left-handed spinor without ambiguity. Our Lagrangian now consists of the Glashow–Salam–Weinberg (GSW) Lagrangian for the electroweak interaction [7–9], plus extra terms in G , H , and N , and subject to the additional constraint

$$g' = \frac{g}{\sqrt{3}}. \quad (19)$$

That is,

$$\mathcal{L} = \mathcal{L}_{\bar{\partial}} + \mathcal{L}_{\text{GSW}} + \mathcal{L}_{\text{extra}} \quad (20)$$

$$\mathcal{L}_{\bar{\partial}} = i\bar{e}_R \bar{\partial} e_R + i\bar{e}_L \bar{\partial} e_L + i\bar{\nu}_e \bar{\partial} \nu_e - \frac{1}{4} F^{\mu\nu} F_{\mu\nu} \quad (21)$$

$$\begin{aligned} \mathcal{L}_{\text{GSW}} = & -g' \left(\bar{e}_R \sigma^\mu e_R + \frac{1}{2} \bar{e}_L \bar{\sigma}^\mu e_L + \frac{1}{2} \bar{\nu}_e \bar{\sigma}^\mu \nu_e \right) X_\mu \\ & + \frac{g}{2} (\bar{\nu}_e \bar{\sigma}^\mu \nu_e - \bar{e}_L \bar{\sigma}^\mu e_L) W_\mu^3 \\ & + \frac{g}{\sqrt{2}} (\bar{e}_L \bar{\sigma}^\mu \nu_e W_\mu + \text{h.c.}) \end{aligned} \quad (22)$$

$$\begin{aligned} \mathcal{L}_{\text{extra}} = & \frac{g}{\sqrt{2}} ([e_R \bar{\sigma}^\mu e_L H_\mu + e_R \bar{\sigma}^\mu \nu_e G_\mu] + \text{h.c.}) \\ & + \bar{\nu}_e \bar{\sigma}^\mu \nu_e N_\mu. \end{aligned} \quad (23)$$

The photon and the Z boson are then constructed via

$$A_\mu = \frac{g' W_\mu^3 + g X_\mu}{(g^2 + g'^2)^{1/2}} \quad (24)$$

$$= X_\mu \cos \theta_W + W_\mu^3 \sin \theta_W \quad (25)$$

$$Z_\mu = \frac{g W_\mu^3 - g' X_\mu}{(g^2 + g'^2)^{1/2}} \quad (26)$$

$$= W_\mu^3 \cos \theta_W - X_\mu \sin \theta_W. \quad (27)$$

The equation

$$\tan \theta_W = \frac{g'}{g} \quad (28)$$

defines the electroweak mixing angle, or Weinberg angle, θ_W , though this is typically measured via the relationship [5, 6]

$$\sin^2 \theta_W = 1 - \frac{m_W^2}{m_Z^2}. \quad (29)$$

In the standard model these expressions coincide, but in the present model we will see in Sec. IIIB that they do not. We will therefore distinguish between the geometric Weinberg angle θ_W^G , defined through (28), and the massive Weinberg angle θ_W^M , defined through (29). For the present model, we find

$$\sin^2 \theta_W^G = 0.25 \text{ (exact)}. \quad (30)$$

We will see that the predicted massive Weinberg angle is slightly smaller, although there is still some discrepancy between this and the measured value [5, 6, 10] of

$$\sin^2 \theta_W^M \text{ (measured)} = 0.222 \ 55(56). \quad (31)$$

III. PARTICLE MASSES

A. Electron mass

In Sec. IIA we postulated the existence of a symmetry breaking mechanism which assigned non-zero values to $\bar{\nu}_e \bar{\sigma}^\mu \nu_e$, and possibly also to $\bar{e}_R \sigma^\mu e_R$, $\bar{e}_L \bar{\sigma}^\mu e_L$, $e_R \bar{\sigma}^\mu e_L$, and $\bar{e}_L \bar{\sigma}^\mu \bar{e}_R$ subject to the requirements of $SU(2)$ symmetry of the electron sector. As the boson fields in the vacuum state must be in equilibrium with the fermion fields, this also implies non-zero values for A_μ , H_μ , $H_\mu^\dagger$, and N_μ in the ground state, along with

$$W_\mu = W_\mu^\dagger = G_\mu = G_\mu^\dagger = Z_\mu = \{0, 0, 0, 0\}. \quad (32)$$

Although the symmetry breaking mechanism determines a preferred orientation on the internal space, it does not select a preferred orientation in space–time, and minimisation of the Lagrangian will result in field configurations satisfying the Euler-Lagrange equations of motion, and

hence periodic in space and/or time. Consequently, averaged over a sufficiently large region,

$$\langle \bar{\phi}_a \bar{\sigma}^\mu \phi_b \rangle = \{0, 0, 0, 0\} \quad \forall \quad a, b \quad (33)$$

$$\langle b_\mu^a \rangle = \{0, 0, 0, 0\} \quad \forall \quad a, \quad (34)$$

except for $\langle \bar{\phi}_3 \bar{\sigma}^\mu \phi_3 \rangle$ which acquires nonzero mean value through our choice of gauge. For brevity let us also introduce the notation

$$\|V_\mu\|^2 = V^\mu V_\mu^\dagger \quad (35)$$

for any 4-vector V , and note that it is the mean ground state magnitudes squared, such as $\langle \|A_\mu\|^2 \rangle$, which are non-zero.

If we now introduce an electron as an excitation above the ground state, it will couple to the vacuum A and H fields according to Lagrangian terms

$$\begin{aligned} \mathcal{L}_{\text{partial}} = & \frac{g}{\sqrt{2}} (e_R \bar{\sigma}^\mu e_L H_\mu + \bar{e}_L \bar{\sigma}^\mu \bar{e}_R H_\mu^\dagger) \\ & - \frac{g}{2} (\bar{e}_R \sigma^\mu e_R + \bar{e}_L \bar{\sigma}^\mu e_L) A_\mu. \end{aligned} \quad (36)$$

Contributions to the path integral which are first order in the boson fields vanish by (34), and to second order we have the interactions

$$\begin{aligned} & \frac{g^2}{4} \bar{e}_L \bar{\sigma}^\mu e_L \bar{e}_L \bar{\sigma}^\nu e_L \langle A_\mu A_\nu \rangle \\ & + \frac{g^2}{2} \bar{e}_L \bar{\sigma}^\mu \bar{e}_R e_R \bar{\sigma}^\nu e_L \langle H_\mu^\dagger H_\nu \rangle \end{aligned} \quad (37)$$

for the propagation of a left-handed electron, and

$$\begin{aligned} & \frac{g^2}{4} \bar{e}_R \sigma^\mu e_R \bar{e}_R \sigma^\nu e_R \langle A_\mu A_\nu \rangle \\ & + \frac{g^2}{2} \bar{e}_R \sigma^\mu \bar{e}_L e_L \sigma^\nu e_R \langle H_\mu^\dagger H_\nu \rangle \end{aligned} \quad (38)$$

for a right-handed electron. We may compare this to the Dirac Lagrangian for an electron in its rest frame (written in terms of Weyl spinors),

$$\mathcal{L}_{\text{Dirac}} = i \bar{e}_R \not{\partial} e_R + i \bar{e}_L \not{\partial} e_L + m_e e_R \sigma^0 e_L + m_e \bar{e}_L \sigma^0 \bar{e}_R, \quad (39)$$

which also gives rise to first as well as second order terms.

For the Dirac Lagrangian, the first order terms are eliminated by the imposition of conservation of charge as an external superselection rule. The second order terms are then

$$m_e^2 \bar{e}_L \sigma^0 \bar{e}_R e_R \sigma^0 \bar{e}_L \quad (40)$$

for the left-handed electron and

$$m_e^2 \bar{e}_R \sigma^0 \bar{e}_L e_L \sigma^0 \bar{e}_R \quad (41)$$

for the right-handed electron respectively. We may identify the Dirac expressions with those in (37) and (38), for which the nonzero mean vacuum expectation values $\langle \|A_\mu\|^2 \rangle$ and $\langle \|H_\mu\|^2 \rangle$ give rise to electron mass.

This should not, however, be taken as implying that $\langle A^\mu A_\mu \rangle = 0$, as the intermediate electron state is undetectable and physically irrelevant to the propagation process. Instead, from the $SU(2)$ symmetry of the electron sector and our choice of basis C_a , we identify

$$\langle \|H_\mu\|^2 \rangle = \langle \|H_\mu^\dagger\|^2 \rangle = \frac{m_e^2}{g^2} \quad (42)$$

$$\langle \|A_\mu\|^2 \rangle = 2 \langle \|H_\mu\|^2 \rangle = \frac{2m_e^2}{g^2} \quad (43)$$

for the vacuum state, where m_e is the observed mass of the electron. Noting from (43) that the background A , H , and $H^\dagger$ fields share a common spatial anisotropy frame, we obtain (40) and (41) directly from (37) and (38) simply by evaluating the Einstein sums in that frame, and find that the rest frame of the electron corresponds directly to the local anisotropy frame of the background A , H , and $H^\dagger$ fields.

There are also higher order interactions which contribute to the electron mass, necessitating small corrections to the above expression for $\langle \|H_\mu\|^2 \rangle$. Fortunately, determining these corrections is straightforward and shall be addressed in Sec. IV C.

B. Boson masses

1. Number density convention

Evaluating Feynman diagrams for interacting particle fields necessitates adopting a convention for interpreting the field strengths as a particle number density. We will take the following approach, which yields mutually consistent interpretations of both the fermion and boson fields:

Let L be unit length. The fermion fields, having units of L^{-3} , may readily be identified with a particle number density. For an arbitrary fermion ϕ_a , the magnitude $\|\bar{\phi}_a \bar{\sigma}^\mu \phi_a\|$ corresponds to the number of particles found in unit volume, where that volume is orthogonal to vector $\bar{\phi}_a \bar{\sigma}^\mu \phi_a$.

For bosons, consider the expression

$$\tilde{g} \bar{\phi}_a \bar{\sigma}^\mu \phi_a b_\mu \quad (44)$$

where $\tilde{g}$ is a coupling constant related to g . The product $\tilde{g} b_\mu$ represents the number of boson interactions occurring for fermions $\bar{\phi}_a \bar{\sigma}^\mu \phi_a$ per unit time, and we may expand this as $\tilde{g} L^3 L^{-3} b_\mu$. The product $\tilde{g} L^3$ then plays the role of the effective volume of one fermion, and $L^{-3} b_\mu$ is the number of boson interactions per unit volume of fermions per unit time. An effective boson number density may be obtained by saturating unit volume of space with fermions, by taking

$$\tilde{g} L^3 \|\bar{\phi}_a \bar{\sigma}^\mu \phi_a\| = 1. \quad (45)$$




FIG. 1: Lowest order diagrams contributing to W and Z boson mass. Grey circles indicate interactions with the background field.

The number of bosons traversing unit volume per unit time is then given by

$$\frac{\tilde{g}L^3 \|\bar{\phi}_a \bar{\sigma}^\mu \phi_a\| b_\mu}{L^3} \quad (46)$$

$$= \frac{b_\mu}{L^3}.$$

2. W , Z , N , and G masses

The primary mechanism responsible for imparting mass to W and Z bosons is postulated to be interactions with the non-zero $\langle \bar{\nu}_e \bar{\sigma}^\mu \nu_e \rangle$ field of the vacuum state. From the standard model we know that the relevant mass terms take the form $m_W^2 W^\dagger W$ and $m_Z^2 Z Z$. Starting from the vacuum state, we introduce a single boson excitation of the desired type. Figure 1 then shows tree diagrams contributing to m_W^2 and m_Z^2 which are first order in the background field.

Evaluating these Feynman diagrams for fixed initial position x gives

$$\int d^4x' W^\dagger(x') [\bar{\nu}_e(x')_{\text{bg}}] \bar{\sigma}^\mu e_L(x') \Delta G_{e_L}(x' - x) \quad (47)$$

$$\times \bar{e}_L(x) \bar{\sigma}^\mu [\nu_e(x)_{\text{bg}}] W(x)$$

and

$$\int d^4x' Z(x') [\bar{\nu}_e(x')_{\text{bg}}] \bar{\sigma}^\mu \nu_e(x') \Delta G_{\nu_e}(x' - x) \quad (48)$$

$$\times \bar{\nu}_e(x) \bar{\sigma}^\mu [\nu_e(x)_{\text{bg}}] Z(x)$$

respectively, where $[\text{bg}]$ denotes a field takes its local vacuum state value. Boson masses then correspond to

$$m_W^2 \equiv \int d^4x' [\bar{\nu}_e(x')_{\text{bg}}] \bar{\sigma}^\mu e_L(x') \Delta G_{e_L}(x' - x) \quad (49)$$

$$\times \bar{e}_L(x) \bar{\sigma}^\mu [\nu_e(x)_{\text{bg}}]$$

$$m_Z^2 \equiv \int d^4x' [\bar{\nu}_e(x')_{\text{bg}}] \bar{\sigma}^\mu \nu_e(x') \Delta G_{\nu_e}(x' - x) \quad (50)$$

$$\times \bar{\nu}_e(x) \bar{\sigma}^\mu [\nu_e(x)_{\text{bg}}].$$




FIG. 2: Consequences of exchanging the intermediary neutrino with one from the background field.

Under integration, $\Delta G_{e_L}(x' - x)$ and $\Delta G_{\nu_e}(x' - x)$ reduce to $\delta^4(x' - x)$, and hence $e_L(x') \Delta G_{e_L}(x' - x) \bar{e}_L(x)$ and $\nu_e(x') \Delta G_{\nu_e}(x' - x) \bar{\nu}_e(x)$ reduce to multiples of $\delta^4(x - x') \sigma^0$. For the former, the multiplication factor is 1 as no other electrons are present in the diagram. For the latter, there are other neutrinos present as a result of the non-zero vacuum expectation value of $\bar{\nu}_e \bar{\sigma}^\mu \nu_e$. However, exchanging the intermediary neutrino with one of them, as in Fig. 2, yields a disconnected diagram, which we interpret as two independent events rather than one particle propagating, because there is no connected pathway to relate the Z boson 4-momenta. Consequently $\nu_e(x') \Delta G_{\nu_e}(x' - x) \bar{\nu}_e(x)$ in (50) also reduces to $\delta^4(x - x') \sigma^0$, and our integrals evaluate in the local rest frame of the background neutrino field as

$$m_W^2 = \frac{g^2 \langle \bar{\nu}_e \bar{\sigma}^0 \nu_e \rangle T}{2} \quad (51)$$

$$m_Z^2 = \frac{2g^2 \langle \bar{\nu}_e \bar{\sigma}^0 \nu_e \rangle T}{3} \quad (52)$$

where T is unit time (distinguished from unit length to aid reinsertion of factors of c in systems of units where these are not 1). This gives again, to first order, $\sin^2 \theta_W^M = 0.25$. Note that the appearance of a constant with value T , unit time, is dependent of our choice of convention for particle number density; however, m_W^2 and m_Z^2 are not, as we shall see in Sec. IV B.

Two families of higher-order diagrams give rise to corrections to this initial result. First, there are radiative corrections to the interactions of Fig. 1. The most significant contribution is electromagnetic, and to one loop this will contribute only to m_W^2 , by modifying the propagator of the intermediary electron (all our propagators so far are for bare particles, so radiative corrections to these diagrams must be made to the particle propagators as well as to the interactions). This correction to m_W^2 is by a factor of

$$1 + \frac{\alpha}{2\pi} + O(\alpha^2) \quad (53)$$

where α is the fine structure constant, with m_Z^2 being

unchanged.

The second correction becomes apparent when, on going to higher orders in g , we also permit multiple interactions with the background fields. For W boson propagation, the W boson may interact with the neutrino fields, and also the intermediate electron may interact with the vacuum H , $H^\dagger$, and photon fields. The intermediate neutrino of Z boson propagation, on the other hand, is nearly massless (see Sec. III C) and it may therefore be inferred that its interactions with the background fields are negligible. The diagrams in Fig. 3 correspond to m_W^4 and m_Z^4 , and we see that again, the W boson mass is augmented relative to that of the Z boson. The correction factor this time is

$$1 + \frac{2m_e^2}{3m_Z^2} + O(m_Z^{-4}). \quad (54)$$

The neutrino interactions similarly correct m_Z , but by a factor proportional to $m_{\nu_e}^2/m_Z^2$, which is safely ignored.

While both of these corrections have the effect of reducing $\sin^2 \theta_W^M$, they are insufficient to bring it down to the observed value of 0.222 55(56). This is unsurprising, as there are known to be further weak boson interactions not included in this model. For example, inclusion of particle generations will give rise to additional terms in the W boson propagator where the intermediate particle is a muon rather than an electron. This will further reduce $\sin^2 \theta_W^M$, as only the W boson participates in flavour-changing weak interactions, and hence the Z boson mass again remains unchanged. It is even possible that couplings to virtual quarks may play a role. That the computed value of $\sin^2 \theta_W^M$ in the partial model presented in this paper is too high is therefore to be expected.

The N and G bosons also couple to $\langle \bar{\nu}_e \bar{\sigma}^\mu \nu_e \rangle$, and hence an N or G boson propagating in the presence of vacuum will exhibit rest mass. Following the same process as above, their masses (to lowest order) are found to be

$$m_N^2 = m_G^2 = \frac{g^2 \langle \bar{\nu}_e \bar{\sigma}^0 \nu_e \rangle T}{2}. \quad (55)$$

In terms of known boson masses, because the N boson interacts only with neutrinos and W bosons, the above corrections are likely to have little effect so we may write

$$m_N^2 \approx 3m_Z^2/4. \quad (56)$$

For the G boson, given its ability to interact with right-handed electrons and the $SU(2)$ symmetry on the ϕ_1, ϕ_2 sector of Φ , it is probably more appropriate to write this as

$$m_G^2 = m_W^2. \quad (57)$$

3. A and H masses

The photon, A , does not couple to $\bar{\nu}_e \bar{\sigma}^\mu \nu_e$ and so does not acquire mass via the mechanism described above. In

addition to the couplings of the standard model, however, the photon also couples to the charged H and G bosons.

We saw in Sec. II A that the bosons coupling to $\bar{\phi}_1 \bar{\sigma}^\mu \phi_3$, $\bar{\phi}_2 \bar{\sigma}^\mu \phi_3$, $\bar{\phi}_3 \bar{\sigma}^\mu \phi_1$, and $\bar{\phi}_3 \bar{\sigma}^\mu \phi_2$ are everywhere zero in the vacuum state, and this corresponds to

$$[W_{\text{bg}}]_\mu = [W_{\text{bg}}^\dagger]_\mu = [G_{\text{bg}}]_\mu = [G_{\text{bg}}^\dagger]_\mu = 0. \quad (58)$$

While in general $[H_{\text{bg}}]_\mu = [H_{\text{bg}}^\dagger]_\mu \neq 0$, this has no effect on photon mass: The A - H coupling in the Lagrangian is

$$\mathcal{L}_{\text{partial}} = \frac{g^2}{2} (H^\mu H_\mu^\dagger A^\nu A_\nu + H^{\dagger\mu} H_\mu A^\nu A_\nu - H^\mu A_\mu H^{\dagger\nu} A_\nu - H^{\dagger\mu} A_\mu H^\nu A_\nu). \quad (59)$$

Expanding in terms of foreground (excitation) fields and background (vacuum expectation value) fields, we find

$$\begin{aligned} \mathcal{L}_{\text{partial}} = \frac{g^2}{2} \bigg[& ([H_{\text{bg}}]^\mu [H_{\text{bg}}^\dagger]_\mu [A_{\text{bg}}]^\nu [A_{\text{bg}}]_\nu \\ & - [H_{\text{bg}}]^\mu [A_{\text{bg}}]_\mu [H_{\text{bg}}^\dagger]^\nu [A_{\text{bg}}]_\nu \\ & + [H_{\text{bg}}]^\mu [H_{\text{bg}}^\dagger]_\mu [A_{\text{fg}}]^\nu [A_{\text{fg}}]_\nu \\ & + [H_{\text{bg}}]^\mu [H_{\text{bg}}^\dagger]_\mu [A_{\text{bg}}]^\nu [A_{\text{fg}}]_\nu \\ & - [H_{\text{bg}}]^\mu [A_{\text{fg}}]_\mu [H_{\text{bg}}^\dagger]^\nu [A_{\text{bg}}]_\nu \\ & - [H_{\text{bg}}]^\mu [A_{\text{bg}}]_\mu [H_{\text{bg}}^\dagger]^\nu [A_{\text{fg}}]_\nu \\ & + [H_{\text{bg}}]^\mu [H_{\text{bg}}^\dagger]_\mu [A_{\text{fg}}]^\nu [A_{\text{fg}}]_\nu \\ & - [H_{\text{bg}}]^\mu [A_{\text{fg}}]_\mu [H_{\text{bg}}^\dagger]^\nu [A_{\text{fg}}]_\nu \bigg) \\ & + (\text{same again with } H \leftrightarrow H^\dagger) \bigg]. \end{aligned} \quad (60)$$

Exploiting (43), all terms save those quadratic in $[A_{\text{fg}}]$ vanish. For these, we note that if the photon acquires a rest mass due to interaction with the background H and $H^\dagger$ fields, then A will share a common rest frame with H and $H^\dagger$. Evaluating in this rest frame, these last terms are also seen to cancel out, so the photon cannot acquire an effective mass from its interaction with H and $H^\dagger$ bosons.

Higher order corrections to the photon propagator also do not give rise to a rest mass. In the standard model, the ability of a photon to generate virtual W and $W^\dagger$ pairs while propagating has no effect on its rest mass, and these couplings are unchanged in the present model. By the $SU(2)$ symmetry of the ϕ_1, ϕ_2 sector in free space, we may conclude that creation of G and $G^\dagger$ pairs also does not give rise to a photon mass, and for H and $H^\dagger$ pairs, the coupling coefficient is different but the form of the interaction is the same. Consequently the photon remains massless.

Finally, the H and $H^\dagger$ bosons are also massless, as $SU(2)$ symmetry of the ϕ_1, ϕ_2 sector in free space guarantees that $m_H = m_{H^\dagger} = (1/\sqrt{2})m_A = 0$.




FIG. 3: Zero-loop diagrams contributing to m_W^4 and m_Z^4 .

C. Neutrino mass

Next we turn to the mass of the electron neutrino, which may also be calculated in this model. A foreground electron neutrino couples to the vacuum N boson field, which will be non-zero as it couples in turn to the vacuum $\bar{\nu}_e \bar{\sigma}^\mu \nu_e$ field.

By $SU(3)$ symmetry, we may anticipate

$$\|[N_{\text{bg}}]_\mu\| = \|[H_{\text{bg}}]_\mu\| = \frac{m_e}{g}. \quad (61)$$

However, this does not confer a mass m_e upon the electron neutrino. Instead, $(g/\sqrt{2})[N_{\text{bg}}]_0$ (in the rest frame of the background field) corresponds to the total mass of all the electron neutrinos in unit volume represented by field $\bar{\nu}_e \bar{\sigma}^\mu \nu_e L^3$. Consequently, the mass a single electron neutrino acquires from interactions with the background N field is equal to

$$\begin{aligned} m_{\nu_e} &= \frac{g}{\sqrt{2}} \frac{m_e}{g} \frac{1}{\langle \|\bar{\nu}_e \bar{\sigma}^\mu \nu_e\| \rangle L^3} \\ &= \frac{\sqrt{2} g^2 m_e T}{3 m_Z^2 L^3}, \end{aligned} \quad (62)$$

and we shall compute the value of g for this model in Sec. IV B. Inserting this value from (81), we find

$$m_{\nu_e} \geq 2.353 \, 64(11) \times 10^{-66} \text{ eV}/c^2, \quad (63)$$

easily compatible with current upper bounds on neutrino mass [10, 11]. This value is given as a lower bound, as it will be supplemented by radiative corrections to the electron neutrino propagator.

IV. BOSON EXCHANGE INTERACTIONS

A. The H interaction

In this section we characterise the interaction consisting of exchange of H and $H^\dagger$ bosons. The existence of these bosons is a major difference between the model presented here and the standard model. However, we shall see that the role of the H bosons, aside from giving rise to electron mass, appears largely insignificant. It is interesting to ask whether an experiment capable of detecting the existence of H and $H^\dagger$ bosons can be devised.

1. The Isolated $H^{(\dagger)}$ boson

First, why do we not detect isolated $H^{(\dagger)}$ bosons? (I will use the notation $H^{(\dagger)}$ to mean “either H or $H^\dagger$ ”.) They should be conspicuous entities, being massless but carrying a charge of ± 2 . The answer lies in the interactions which generate them. A single H^{--} boson may be generated by the interaction

$$e_L + e_R \longrightarrow H^{--}. \quad (64)$$

However, this interaction destroys a left-handed electron and a right-handed electron. By implication, the H boson created has mass $2m_e$, and so is clearly virtual. Interestingly, it also has lepton number 2. Emission of a non-virtual H boson can only be achieved if additional virtual particles are also near-simultaneously created which share a combined mass deficit of $2m_e$. The only realistic

candidate is creation of an H^{++} boson, as in the example

$$\begin{aligned} e_L &\longrightarrow \bar{e}_R + H^{--} \\ \bar{e}_R + H^{--} &\longrightarrow e_L + H^{--} + H^{++}. \end{aligned} \quad (65)$$

Similarly, the H and $H^\dagger$ bosons may also only be absorbed in an appropriate pairwise fashion. Considered collectively, the H , $H^\dagger$ pair has zero charge, zero mass, and zero lepton number. Like a pair of photons, they may carry a spin of zero or ± 2 . Indeed, it would seem that distinguishing between emission or absorption of a pair of photons and emission or absorption of a pair of H and $H^\dagger$ bosons will be very difficult indeed. This is, of course, only to be anticipated as the H and $H^\dagger$ bosons share an $SU(2)$ symmetry sector with the photon, and that symmetry is unbroken.

2. $H^{(\dagger)}$ boson exchange

So what are the consequences of $H^{(\dagger)}$ exchange? Is there any prospect for detecting this interaction as a macroscopic force? Once again, we may obtain our answer via the unbroken $SU(2)$ symmetry of the ϕ_1, ϕ_2 sector. We will take as a basis for the $su(2)$ Lie algebra the coupling of the photon A_μ , and the couplings of the linear combinations of $H^{(\dagger)}$ bosons

$$H_\mu^1 = \frac{1}{\sqrt{2}} (H_\mu + H_\mu^\dagger) \quad (66)$$

$$H_\mu^2 = \frac{1}{\sqrt{2}} (iH_\mu^\dagger - iH_\mu). \quad (67)$$

The electromagnetic interaction is mediated by the photon, which couples with strengths $+e$ and $-e$ to positrons and electrons respectively. We therefore say that, relative to the electromagnetic interaction, these particles have charges ± 1 . What are the charges of electrons with regards to the H^1 and H^2 bosons?

To resolve this question, we identify the $SU(2)$ mixings of ϕ_1 and ϕ_2 (e_L and $\bar{e}_R$) which diagonalise the H^1 boson. They are

$$h_1^1 = \pm \frac{1}{\sqrt{2}} (e_L + \bar{e}_R) \quad (68)$$

$$h_2^1 = \pm \frac{1}{\sqrt{2}} (e_L - \bar{e}_R) \quad (69)$$

and they carry charges relative to the H^1 -boson-mediated interaction of 1 and -1 respectively. From this we see that electrons constitute an equal mixing of constituents with H^1 -charges $+1$ and -1 . Consequently the net H^1 -charge of the electron or the positron is zero, and when exchanged between electrons or positrons the H^1 boson yields no net force. The argument may be repeated for interactions mediated by the H^2 boson.

An extension of the same argument holds for H -mediated interactions between the W and G bosons, and




FIG. 4: Electromagnetic interaction mediated by virtual photon exchange. Labelling e rather than e_L or e_R indicates that the electrons may be of either spin.

the H -mediated interactions of the photon cancel by symmetry as H , $H^\dagger$, and A form an orthonormal basis of the unbroken $SU(2)$ symmetric subspace. Neither the N nor the Z bosons interact with the $H^{(\dagger)}$ at all, and we are left to conclude that the $H^{(\dagger)}$ bosons mediate no classical force.

3. Prospects for detection of $H^{(\dagger)}$ bosons

We have seen that the $H^{(\dagger)}$ boson is very difficult to detect. It is never emitted or absorbed singly, $H^{(\dagger)}$ boson exchange mediates no classical force, and pairs of $H^{(\dagger)}$ bosons have zero net charge, zero mass, and a combined spin of 0 or ± 2 , like pairs of photons. Proof or refutation of their existence is therefore rather difficult. One approach to test for the existence of the H boson would be to exploit its existence as an alternative virtual intermediary for photon-mediated interactions. For example, compare

$$e_L + \bar{e}_R + \bar{e}_L \longrightarrow \bar{e}_R + A \longrightarrow \bar{e}_R \quad (70)$$

$$e_L + \bar{e}_R + \bar{e}_L \longrightarrow e_L + H^{++} \longrightarrow \bar{e}_R. \quad (71)$$

This alternative pathway, if it exists, should result in an increase in the apparent rate of electron-positron annihilation when a second positron of opposite spin absorbs the gamma ray which is produced. The author is unaware of any experimental observations which verify or refute the existence of $H^{(\dagger)}$ bosons.

B. Electromagnetism

We turn now to the electromagnetic interaction, mediated by virtual photon exchange (Fig. 4). The presence of a non-zero vacuum expectation value of the photon field significantly modifies this interaction, and we will find that the value of g coinciding with observed behaviour differs from its usual value in quantum field theory of $g^2 = e^2/(\epsilon_0 \sin^2 \theta_W)$.

The origin of this correction lies in the indistinguishability of photons, and consequently we cannot uniquely identify the photon exchanged between the two electrons

in Fig. 4. Instead, all we can say is that one electron emits a photon at point x , and the other absorbs one at point x' . We require that when we average over sufficiently many such interactions the 4-momentum of the background field remains unchanged, so we may say that emission of a photon at x and absorption at x' has resulted in a transfer of momentum from one electron to the other.

The numerical factor associated with Fig. 4 for a pair of left-handed electrons interacting with the photon field at x and x' is

$$\begin{aligned} & \iint d^4x d^4x' \frac{g^2}{2} \bar{e}_L(x) \bar{\sigma}^\mu e_L(x) \{ [A_{bg}]_\mu(x) + [A_{fg}]_\mu(x) \} \\ & \times \Delta_A(x - x') \bar{e}_L(x') \bar{\sigma}^\mu e_L(x') \{ [A_{bg}]_\mu(x') + [A_{fg}]_\mu(x') \}. \end{aligned} \quad (72)$$

The four permutations of the external legs, and the symmetry factor $1/2$ from exchange of the interaction vertices, have been combined with the boson couplings $(g/2)^2$ to give the overall leading factor of $g^2/2$.

Terms involving $[A_{bg}]_\mu$ at both x and x' are unchanged from electron propagation in free space and so do not correspond to momentum transfer. Of the remaining three terms, two involve both the background field and the foreground field, and one involves the foreground field only. For sufficiently large separations $[A_{bg}]_0(x) \gg [A_{fg}]_0(x)$ in the rest frame of the absorbing electron, and we may neglect the term quadratic in the foreground field. Substituting (42-43) into the remaining two terms, taking

$$\Delta_A(r, t) = \frac{e^{i\theta(r, t)}}{4\pi r^2}, \quad (73)$$

and employing the number density convention of Sec. III B 1 to evaluate the foreground fields, we obtain a value of

$$F = \frac{\sqrt{2} g m_e T}{4\pi r^2} e^{i\theta(r, t)} \quad (74)$$

where T represents unit time. On the light cone this must correspond to a classical interaction strength $F = e^2/(4\pi r^2 \epsilon_0)$, and consequently

$$g = \frac{e^2}{\sqrt{2} \epsilon_0 m_e T}. \quad (75)$$

This expression is valid for $[A_{bg}]_0(x) \gg [A_{fg}]_0(x)$ in the instantaneous rest frame of the absorbing electron, corresponding to $r \gg r_{\text{crit}}$ where

$$\begin{aligned} r_{\text{crit}} &= \sqrt{\frac{g^2 T}{8\pi \sqrt{2} m_e}} + O(g^2) \\ &\sim 5.3 \times 10^{-25} \text{ m}. \end{aligned} \quad (76)$$

It might give the reader momentary pause to note that g is quadratic in the electron charge e . However, this

does not equate to the electromagnetic interaction being quadratic in e because the background photon field introduces an additional factor of $\sqrt{2} m_e/g$. Transfer of momentum via single photon exchange is therefore, to lowest order, linear in g and hence is quadratic in e as is expected. Indeed, the value of g is chosen such that despite the presence of a background field, the factor associated with single photon exchange is unchanged from its usual value in quantum electrodynamics. At present, this is true only if radiative corrections are ignored, but it is easy to determine the necessary small correction to g such that this correspondence holds for interactions calculated to all orders, and this is the subject of Sec. IV C.

Another important point is that the numerical value of g is dependent upon the number density convention adopted. Changing the number convention means that the field distribution associated with a single foreground electron or photon in (72) changes, but the background field is unaffected. Consequently, identifying the relevant terms of (72) with the classical interaction strength will yield a different value of g . It is an important principle that any measurable physical property should be independent of our choice of number density convention. For example, if we write

$$\int dV \bar{e}_L \bar{\sigma}^0 e_L = k N_{e_L} \quad (77)$$

then the convention of Sec. III B 1 corresponds to $k = 1$. Adopting an alternate value of k gives $g \rightarrow g/k$, but because $[A_{bg}]_\mu$ and $[H_{bg}^{(\dagger)}]_\mu$ remain unchanged, we must also have $m_e \rightarrow m_e/k$. The electron mass interaction terms of $\mathcal{L}$ then become

$$\frac{m_e}{k} \bar{e}_L \bar{\sigma}^\mu \bar{e}_R + \frac{m_e}{k} e_R \bar{\sigma}^\mu e_L \quad (78)$$

but the measured value of the electron mass remains unchanged due to the appearance of k in (77) when integrating these terms. Similarly the measured boson and neutrino masses are also unaffected. Note that our expression for neutrino mass depends on the assumption that

$$\left\| [\bar{\nu}_e \bar{\sigma}^\mu \nu_e]_{\text{fg}} \right\| \ll \left\| [\bar{\nu}_e \bar{\sigma}^\mu \nu_e]_{\text{bg}} \right\|.$$

If this assumption fails to hold the expression for the measured neutrino mass m_{ν_e} will change, but the calculated value will remain the same.

The calculated value of r_{crit} , on the other hand, will vary with the definition of number density. This is unsurprising, as it is defined as the radius at which the foreground photon field of *one* electron becomes comparable to the background fields, and this is dependent upon the definition of the field excitation corresponding to one electron. When giving a value for r_{crit} , it is therefore necessary to specify also the number density convention to which it corresponds. We will use only the $k = 1$ convention in this paper.




FIG. 5: Feynman diagrams for the electron mass interaction.

C. Radiative corrections

In the previous section, we calibrated the order g^2 electromagnetic interaction to be compatible with classical electrodynamics. This section deals briefly with the subject of radiative corrections to the electromagnetic interaction, and also to the electron mass.

First, note that the factors associated with an internal photon line, although more complicated in origin, are now no different to those in normal quantum electrodynamics due to our calibration of g in Sec. IV B. Contributions to higher-order interactions due to internal photon lines are therefore unchanged.

Second, recall that the $H^{(\dagger)}$ bosons mediate no force, and hence carry no net momentum, so diagrams involving $H^{(\dagger)}$ boson exchange make zero net contribution to radiative corrections.

Third, there are no background fields for the W , Z , or G bosons. The W and Z boson radiative corrections are therefore unchanged from the GSW model. G bosons may also be ignored in curved space-time, for reasons to be explained in Sec. V B.

There will be changes to higher-order radiative correction terms, but only where these involve neutrino propagators (or N boson propagators, if these are assumed to exist). Such corrections are anticipated to lie at the present limits of theoretical and experimental accuracy, and consequently existing results for the gyromagnetic ratio of the electron continue to hold, at least to the level where the weak interaction becomes significant (in the standard model, $0.2564(45) \times 10^{-10} a_e$, where a_e is the electron magnetic moment anomaly [5, 6]).

To compute the value of g after radiative corrections, first we must evaluate their effects on the background fields. As in Sec. III A, we consider interactions giving the electron mass. The zero order Feynman diagrams are shown in Fig. 5, and the first order radiative corrections are shown in Fig. 6. Four of these figures have positive sign and two have negative sign. The net effect is that radiative corrections to the electron mass go as $(1 + a_e)$ per coupling to the background field. Consequently we

must amend our mean square background field values to

$$\langle \|H_\mu\|^2 \rangle = \langle \|H_\mu^{(\dagger)}\|^2 \rangle = \frac{m_e^2}{g^2(1 + a_e)^2} \quad (79)$$

$$\langle \|A_\mu\|^2 \rangle = 2\langle \|H_\mu\|^2 \rangle = \frac{2m_e^2}{g^2(1 + a_e)^2}. \quad (80)$$

The value of g is again calculated by requiring that the electromagnetic interaction strength obtained is in agreement with observation, only now the interaction vertices are associated with factors of $g(1 + a_e)$ representing the electromagnetic interaction to all orders, and we adopt the new values (79-80) for the rest frame background fields. The result (for this number density convention) is

$$\begin{aligned} g &= \frac{e^2}{\sqrt{2}\varepsilon_0 m_e T(1 + a_e)c^3} \\ &= 8.342\,622\,85(59) \times 10^{-23}. \end{aligned} \quad (81)$$

V. GRAVITATION

Finally, we are in a position to study the interaction mediated by the $G^{(\dagger)}$ bosons, which will turn out to be a proxy for the gravitational field. Our choice of gauge at the end of Sec. II A means that the vacuum expectation values $\langle G_\mu^{(\dagger)} \rangle$ are zero. However, a foreground lepton may still directly generate a $G^{(\dagger)}$ field. What is worse, such a field appears to produce a form of right-handed weak interaction. In this section we will examine the interactions of the $G^{(\dagger)}$ fields, and see how a choice of gauge on part of the combined internal and external symmetry group $GL(3, \mathbb{R}) \otimes P_{3,1}$ enables us to eliminate all relevant interactions involving $G^{(\dagger)}$ bosons.

As well as interacting with leptons, $G^{(\dagger)}$ bosons may interact with other bosons. In the virtual fields surrounding an electron, we may identify two distinct domains: At small r , the $G^{(\dagger)}$ fields are comparable in magnitude to the fields of the other bosons, and processes generating and eliminating pairs of $G^{(\dagger)}$ bosons exist in a state approaching equilibrium. However, at larger radii an asymmetry develops. The existence of non-zero vacuum expectation values for certain other bosons but not for the $G^{(\dagger)}$ particles drives $G^{(\dagger)}$ production more rapidly than $G^{(\dagger)}$ annihilation. Because of the requirement that they have a non-zero vacuum expectation value, the only boson fields capable of participating significantly in $G^{(\dagger)}$ creation at large r are those with non-zero local vacuum expectation values— A , $H^{(\dagger)}$, and N .

A. $G^{(\dagger)}$ field of a neutral body

1. Radial dependence

Consider the boson fields surrounding a spherically symmetric, irrotational collection of an equal number of




FIG. 6: Radiative corrections to the mass interaction of the left-handed electron. Diagrams (i)-(iv) have positive sign and diagrams (v)-(vi) have negative sign. Diagrams for the right-handed electron are equivalent.

positrons and electrons. Such a collection will have zero net charge, and will serve as our prototype for a massive, electrically neutral body. Assuming such a body contains both left- and right-handed electrons and positrons, it will generate virtual boson fields of W and $W^\dagger$ bosons, Z bosons, H and $H^\dagger$ bosons, and G and $G^\dagger$ bosons. Of these virtual boson fields, only the $H^{(\dagger)}$ fields have a non-zero vacuum expectation value. We therefore use them to distinguish the two domains in which qualitatively different physics apply: Small r , where the virtual $H^{(\dagger)}$ fields are much greater than the vacuum expectation values, and large r , where the virtual $H^{(\dagger)}$ fields are much smaller than the local vacuum field. At small r , where the background fields are negligible, the $H^{(\dagger)}$ boson density in the neutral body's rest frame is given by

$$H_\mu^{(\dagger)} = \left\{ \frac{H(0)}{4\pi r^2}, 0, 0, 0 \right\}. \quad (82)$$

$$H(0) = \frac{gMT}{\sqrt{2}m_e}. \quad (83)$$

The radial component is zero because, under Feynman propagators, both advanced and retarded fields are present, with incoming and outgoing radial components respectively, which sum to zero.

Equating the virtual boson field (82) with the background field (79) gives the radius at which the transition between the two regimes takes place for the $H^{(\dagger)}$ fields of the neutral object:

$$r_{\text{crit}}^H = \sqrt{\frac{g^2 MT}{8\pi m_e^2}} \quad (84)$$

$$\sim \sqrt{M/m_e} \times 10^{-24} \text{ m}. \quad (85)$$

For simplicity we will treat the object as a point, having radius $r_{\text{object}} \ll r_{\text{crit}}^H$.

Within the low- r domain, couplings between the $G^{(\dagger)}$ fields and all virtual boson fields are significant, and the fields exist in a state approaching equilibrium. Consequently we may also write the $G^{(\dagger)}$ boson fields in the form

$$G_\mu^{(\dagger)} = \left\{ \frac{G(0)}{4\pi r^2}, 0, 0, 0 \right\} \quad (86)$$

$$G(0) = H(0) = \frac{gMT}{\sqrt{2}m_e}, \quad (87)$$

satisfying $G_\mu^{(\dagger)} = H_\mu^{(\dagger)}$ for $r \ll r_{\text{crit}}^H$.

For $r \gg r_{\text{crit}}^H$, a different regime predominates. Here, $H_\mu^{(\dagger)}$ is dominated by the background field, and is much larger than any of the other virtual boson fields. As a result of our choice of gauge in Sec. II A, which prohibits formation of $G^{(\dagger)}$ pairs in the vacuum state, conversion of $H^{(\dagger)}$ bosons into $G^{(\dagger)}$ bosons is dependent upon the presence of the foreground virtual boson field, and this constrains the maximum attainable values of $G^{(\dagger)}$. However, as we shall see, the presence of the background fields

catalyses the interaction, favouring the process

$$H + H^\dagger \longrightarrow G + G^\dagger \quad (88)$$

over its inverse, to the extent that for $r \gg r_{\text{crit}}^H$ the reverse process is negligible. For convenience we shall assume a sharp transition between the $r \ll r_{\text{crit}}^H$ and $r \gg r_{\text{crit}}^H$ regimes occurs at r_{crit}^H , an assumption which introduces negligible error provided r is sufficiently larger than r_{crit}^H .

If we assume that the net rate of interaction (88) is negligible at $r < r_{\text{crit}}^H$, the $H^{(\dagger)}$ field in the neutral body's rest frame at $r = r_{\text{crit}}^H$ is given by (82). Within this radius, the field strength is governed solely by the inverse square law. At $r > r_{\text{crit}}^H$, two independent processes are responsible for decrease of the $H^{(\dagger)}$ fields with r : Ongoing spatial dispersion according to the inverse square law, and also interaction (88). We may thus write the zero component of $H^{(\dagger)}$ field strength for $r > r_{\text{crit}}^H$ as

$$H_0^{(\dagger)}(r) = \lim_{\Delta r \rightarrow 0} \left[\frac{gMT}{4\sqrt{2}\pi(r_{\text{crit}}^H)^2 m_e} \times \left(1 - \frac{2\Delta r}{r}\right)^{(r-r_{\text{crit}}^H)/\Delta r} (1 - \Delta)^{(r-r_{\text{crit}}^H)/\Delta r} \right] \quad (89)$$

where the first infinitesimal corresponds to further application of the inverse square law, and the second corresponds to interaction (88) where Δ is a function of Δr still to be determined. Similarly, while (88) diminishes the $H^{(\dagger)}$ fields, it causes the $G^{(\dagger)}$ fields to increase:

$$G_0^{(\dagger)}(r) = \lim_{\Delta r \rightarrow 0} \left[\frac{gMT}{4\sqrt{2}\pi(r_{\text{crit}}^H)^2 m_e} \times \left(1 - \frac{2\Delta r}{r}\right)^{(r-r_{\text{crit}}^H)/\Delta r} (1 + \Delta)^{(r-r_{\text{crit}}^H)/\Delta r} \right]. \quad (90)$$

2. Evaluating Δ

To evaluate Δ , we must consider the Feynman diagrams for (88) occurring in the $H^{(\dagger)}$ fields surrounding the neutral mass. As we are interested in the total rate of interaction, rather than the scattering cross-section, the most direct way to calculate this is by employing disjoint Feynman diagrams in which the entirety of the source is explicitly represented. Emission of the virtual $H^{(\dagger)}$ fields in the presence of the background fields may be represented by Fig. 7. In calculating Δ we are not especially interested in the factor associated with this diagram, as this is already incorporated in the source term in (89),

$$\frac{gMT}{4\sqrt{2}\pi(r_{\text{crit}}^H)^2 m_e}.$$

We are, however, interested in the degeneracy of the $H^{(\dagger)}$ bosons which are available at radius r . Inverse square




FIG. 7: Emission of (i) an H boson and (ii) an $H^\dagger$ boson by a neutral source of mass M in the presence of locally nonzero background H and $H^\dagger$ fields. The absence of a spin on e and $\bar{e}$ indicates that the diagrams are to be summed over electron spin. As the source is neutral, there are $M/(2m_e)$ electrons and $M/(2m_e)$ positrons in each of diagrams (i) and (ii) respectively.




FIG. 8: Feynman diagram for $H + H^\dagger \rightarrow G + G^\dagger$.

law effects are incorporated into the relevant propagators, and are again realised separately in the other terms of (89), so for computing Δ we are only interested in the number of diagrams delivering bosons to radius r . For the foreground H and $H^\dagger$ bosons, there are $N/2$ diagrams apiece, where $N = M/m_e$ is the number of particles in the electrically neutral massive body. These $N/2$ diagrams corresponding to emission of an H or $H^\dagger$ boson by each of the $N/2$ different electrons (for H) or positrons (for $H^\dagger$) in the source.

For the background fields, by our number convention of Sec. III B 1, there are $[H_{\text{bg}}^{(\dagger)}]_\mu \Delta r^\mu$ background bosons of each type $H^{(\dagger)}$ for every $N/2$ foreground bosons. We can probe $H^{(\dagger)}$ field strength by introducing a single electron or positron at radius r which is stationary in the desired rest frame, and seeing how many bosons it interacts with in the time interval required for the fields to cross radial distance Δr . As the $H^{(\dagger)}$ fields are massless, it follows that the four-vector $\Delta r^\mu = \{\Delta r, 0, 0, 0\}$, explaining the use of the symbol Δr for this four-vector as well as for the spatial distance. Consequently,

$$[H_{\text{bg}}^{(\dagger)}]_\mu \Delta r^\mu = [H_{\text{bg}}^{(\dagger)}]_0 \Delta r. \quad (91)$$

To lowest order, the interaction giving rise to the $G^{(\dagger)}$ fields is shown in Fig. 8. This interaction could in theory take place between any of the H and $H^\dagger$ bosons present

at radius r , foreground or background. However, the choice of gauge in Sec. II A provides that in the absence of any foreground fields, $G_\mu^{(\dagger)}$ is everywhere $\{0, 0, 0, 0\}$. The extension of this choice of gauge to the presence of foreground fields similarly suppresses formation of $G^{(\dagger)}$ bosons, such that it is as if only the foreground bosons are able to participate in the interaction. Similarly, in (89), Δ is also defined acting only on the foreground fields. We may think of Δ^2 as the probability amplitude that a given pair of foreground bosons H and $H^\dagger$ will interact to form a pair G and $G^\dagger$. Naïvely, Fig. 8 might be associated with the factor

$$p = \int_{r_{\text{crit}}^H}^{r_{\text{crit}}^H + \Delta r} dr \frac{g^2}{4} (H^{\dagger\mu} H^\nu G_\mu G_\nu^\dagger + H^\mu H^{\dagger\nu} G_\mu^\dagger G_\nu - H^\mu G^{\dagger\nu} G_\mu H_\nu^\dagger - G^\mu H^{\dagger\nu} H_\mu G_\nu^\dagger - H^\mu H^{\dagger\nu} G_\mu G_\nu^\dagger - H^{\dagger\mu} H^\nu G_\mu^\dagger G_\nu + H^\mu G^{\dagger\nu} H_\mu^\dagger G_\nu + H^{\dagger\mu} G^\nu H_\mu G_\nu^\dagger) \quad (92)$$

but this does not take into account the effect of the choice of gauge restricting the interaction to only the foreground bosons. To include this effect, we multiply (92) by a factor representing the fraction of interactions (88) which take place with only foreground $H^{(\dagger)}$ bosons. Diagrammatically, this factor is shown in Fig. 9.

Assuming $\|[H_{\text{bg}}^{(\dagger)}]\| \gg \|[H_{\text{fg}}^{(\dagger)}]\|$, we can neglect all terms involving foreground fields in the denominator of Fig. 9, and insert (80) and (79) to obtain

$$I = \frac{(N/2)^2}{\frac{1}{2}[H_{\text{bg}}]_0[H_{\text{bg}}^\dagger]_0\Delta r^2} = \frac{M^2 g^2 (1 + a_e)^2}{2m_e^4 \Delta r^2}. \quad (93)$$

The extra factor of a half in the denominator arises because attaching the interaction of Fig. 8 to the back-

$$I = \frac{\text{Diagram 1}}{\text{Diagram 2} + \text{Diagram 3} + \text{Diagram 4} + \text{Diagram 5}}$$

The figure illustrates the fraction of $H + H^\dagger \rightarrow G + G^\dagger$ interactions occurring on the foreground field alone. The numerator represents the interaction where the foreground field alone contributes to the emission of H and $H^\dagger$ bosons. The denominator represents the total interaction, including contributions from the background field. The diagrams show the interaction of $M/(2m_b)$ electrons and $M/(2m_b)$ positrons with H and $H^\dagger$ bosons, leading to the production of G and $G^\dagger$ particles.

FIG. 9: Fraction of $H + H^\dagger \rightarrow G + G^\dagger$ interactions which take place on the foreground field alone, represented as a ratio of Feynman diagrams. The portions of the diagrams enclosed in boxes correspond to emission of the H and $H^\dagger$ bosons by the source. This interaction is accounted for elsewhere, and these portions of the diagram are included only to assist in bookkeeping, being indicative of the degeneracy factors of the foreground and background $H^{(\dagger)}$ bosons.




FIG. 10: Subdiagram corresponding to spontaneous vacuum generation of $GG^\dagger$ pairs.

ground fields removes half of the diagrams in the denominator of I . In the numerator we have $[H_{\text{bg}}]_0 \Delta r$ sources of type H and $[H_{\text{bg}}^\dagger]_0 \Delta r$ sources of type $H^\dagger$ at the origin, and approximately the same number again at radius r (for $r \gg r_{\text{crit}}^H$). These may then be connected in pairs in any manner desired, including where the paired H and $H^\dagger$ are both at r , or both at the origin. In the denominator, however, only half of these diagrams are relevant to the problem at hand. The other half include Fig. 10 as a subdiagram, and so deal with spontaneous generation of $GG^\dagger$ pairs from the vacuum, rather than conversion of $HH^\dagger$ pairs to $GG^\dagger$ pairs as they propagate from r to $r + \Delta r$.

Now that we have obtained the degeneracy factor I , we wish to compute Δ in (89-90), which may be interpreted as a reaction rate per boson. Considering both Figs. 8 and 9, we can write

$$\begin{aligned} \Delta^2 &= -I p_1 = -\frac{M^2 g^2 (1 + a_e)^2 p_1}{2m_e^4 \Delta r^2} \\ &= \frac{M^2 g^4 (1 + a_e)^2}{4m_e^4 \Delta r^2} \end{aligned} \quad (94)$$

where p_1 is Fig. 8 evaluated with only a single H boson and a single $H^\dagger$ boson. Note that as this corresponds to dividing Fig. 8 by the $H^{(\dagger)}$ boson degeneracy, the contribution of pre-existing $G^{(\dagger)}$ fields to p_1 is negligible at $r > r_{\text{crit}}^H$, where $\|G_\mu^{(\dagger)}\| \ll \|H_\mu^{(\dagger)}\|$. The factor p_1 in (94) has therefore been evaluated as if the only $G^{(\dagger)}$ bosons present are those created by the interaction shown. In particular the final two terms in (92) are negligible, as these only affect the propagation of existing $G^{(\dagger)}$ and $H^{(\dagger)}$ excitations, without coupling any new excitations from the $H^{(\dagger)}$ fields into the $G^{(\dagger)}$ fields.

By our choice of sign in (89) and (90), we now have

$$\Delta = \frac{M g^2 (1 + a_e)}{2m_e^2 \Delta r}. \quad (95)$$

In the limit $r \gg r_{\text{crit}}^H$ we may approximate $r_{\text{crit}}^H \rightarrow 0$,

giving

$$H_0^{(\dagger)}(r) = \lim_{\Delta r \rightarrow 0} \left[H(0) \left(1 - \frac{2\Delta r}{r} \right)^{r/\Delta r} \times \left(1 - \frac{M g^2 (1 + a_e)}{2m_e^2 \Delta r} \right)^{r/\Delta r} \right] \quad (96)$$

$$G_0^{(\dagger)}(r) = \lim_{\Delta r \rightarrow 0} \left[G(0) \left(1 - \frac{2\Delta r}{r} \right)^{r/\Delta r} \times \left(1 + \frac{M g^2 (1 + a_e)}{2m_e^2 \Delta r} \right)^{r/\Delta r} \right], \quad (97)$$

and provided we can simultaneously satisfy

$$r \gg \Delta r \gg \frac{M g^2 (1 + a_e)}{2m_e^2}, \quad (98)$$

which is possible for

$$r \gg \frac{M g^2 (1 + a_e)}{2m_e^2}, \quad (99)$$

we may further approximate this as

$$H_0^{(\dagger)}(r) \approx \frac{gMT}{4\sqrt{2}\pi r^2 m_e} \exp \left[-\frac{M g^2 (1 + a_e)}{2m_e^2 r} \right] \quad (100)$$

$$G_0^{(\dagger)}(r) \approx \frac{gMT}{4\sqrt{2}\pi r^2 m_e} \exp \left[\frac{M g^2 (1 + a_e)}{2m_e^2 r} \right]. \quad (101)$$

B. Elimination of the $G^{(\dagger)}$ fields

Now that we have calculated the $G^{(\dagger)}$ fields as a function of r , it is time to eliminate them. We will do this by promoting the global Lorentz invariance of (1) to local Lorentz invariance, and choosing a gauge on the new gauge group $GL(3, \mathbb{R}) \otimes SO(3, 1)^{\uparrow+}$.

A system which is locally Lorentz invariant may be described equally well in any unitary co-ordinate system—that is, one in which the magnitude of the volume form $\sqrt{|g|} d^4x$ is everywhere normalised to one. Promoting Lorentz invariance to a local symmetry thus allows us to adopt a co-ordinate frame on our Minkowski manifold in which the volume form is conserved, but the 4-vectors of the co-ordinate frame need not individually be normalised to one when expressed in terms of Cartesian co-ordinates on Minkowski space-time. The frame may then give rise to a non-trivial metric.

We wish to eliminate any evidence of the $G^{(\dagger)}$ bosons, and we do this by redefining the covariant derivative, adopting a co-ordinate frame satisfying

$$\begin{aligned} \mathcal{D}_\mu &= \partial_\mu - \frac{ig}{\sqrt{2}} \sum_{a=1}^9 b_\mu^a C_a \\ &\downarrow \\ \mathcal{D}'_\mu &= \nabla_\mu - \frac{ig}{\sqrt{2}} \sum_{a \in \{1, 2, 4, 5, 6, 8, 9\}} b_\mu^a C_a \end{aligned} \quad (102)$$

$$\mathcal{D}'_\mu \Phi = \mathcal{D}_\mu \Phi \quad \forall \quad \Phi$$

where the first line of (102) is evaluated in Minkowski space-time, and the second line is evaluated in a locally Lorentz transformed co-ordinate frame. Eq. (102) transforms both the internal and external connections, but leaves the dynamics of the spinor fields unchanged, and so defines a choice of gauge on $GL(3, \mathbb{R}) \otimes SO(3, 1)^{\dagger+}$. While there is no guarantee that this equivalence may be solved in general, appropriate solutions do exist for the spherically symmetric example of Sec. V A. These solutions are all equivalent up to a local change of basis, so we need only consider a typical example of the class.

The defining feature of co-ordinate systems satisfying (102) is that the G and $G^\dagger$ bosons do not participate in any interactions. For our spherically symmetric system, their local spatial profile will therefore be purely inverse square. We see this may be satisfied by defining the new, primed co-ordinate system

$$r' = r \exp \left[-\frac{Mg^2(1+a_e)}{4m_e^2 r} \right] \quad (103)$$

$$\theta' = \theta \quad (104)$$

$$\phi' = \phi \quad (105)$$

$$t' = t \exp \left[\frac{Mg^2(1+a_e)}{4m_e^2 r} \right] \quad (106)$$

where we have chosen t' to conserve the volume form $\sqrt{|g|}d^4x$. The resulting co-ordinate frame is well-defined everywhere save at the origin, where it is impossible to smoothly continue the frame without the radial component going to zero and the time component becoming infinite.

To evaluate the metric for the new co-ordinate frame, we compute

$$dr \approx dr' \exp \left[\frac{Mg^2(1+a_e)}{4m_e^2 r'} \right] \quad (107)$$

$$d\theta = d\theta' \quad (108)$$

$$d\phi = d\phi' \quad (109)$$

$$dt \approx dt' \exp \left[-\frac{Mg^2(1+a_e)}{4m_e^2 r'} \right] \quad (110)$$

where the approximations hold in limit (99). This yields

$$\begin{aligned} ds^2 \approx & \exp \left[-\frac{Mg^2(1+a_e)}{2m_e^2 r'} \right] dt'^2 \\ & + \exp \left[\frac{Mg^2(1+a_e)}{2m_e^2 r'} \right] dr'^2 \\ & + r'^2 (d\theta'^2 + \sin^2 \theta' d\phi'^2), \end{aligned} \quad (111)$$

again in the limit (99). We recognise this as a form of the Schwarzschild metric, with

$$G = \frac{g^2(1+a_e)}{4m_e^2}. \quad (112)$$

This is the metric of our locally Lorentz transformed co-ordinate frame on Minkowski space, but the same metric also describes a co-ordinate frame with each vector independently normalised to 1 on a curved space-time. The two situations are physically indistinguishable, and so the dynamics of all particles except the $G^{(\dagger)}$ bosons in curved space-time are equally well described by the dynamics of all particles including the $G^{(\dagger)}$ bosons on Minkowski space-time. The theory is renormalisable on Minkowski space-time (assuming an appropriate symmetry breaking mechanism), and the use of a gauging process to relate the apparent space-time curvature to matter fields avoids the need for the direct coupling prohibited by the Coleman–Mandulak theorem [12]. What this does suggest, however, is that it may be impossible to construct an equivalent renormalisable theory living purely on the curved space-time.

Given (112), limit (99) reduces to

$$r \ggg 2GM, \quad (113)$$

and in this limit the metric may then be written as

$$\begin{aligned} ds^2 \approx & \left(1 - \frac{2GM}{r'} \right) dt'^2 + \left(1 - \frac{2GM}{r'} \right)^{-1} dr'^2 \\ & + r'^2 (d\theta'^2 + \sin^2 \theta' d\phi'^2). \end{aligned} \quad (114)$$

How accurate is this value of G ? Substituting (81) for g and reinserting the necessary factors of c and $\hbar$, we compute

$$\begin{aligned} G &= \frac{e^4(1+a_e)}{8\varepsilon_0^2 m_e^4 T^2} \frac{\hbar}{c^5} \\ &= 6.652\,336\,1(15) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2} \end{aligned} \quad (115)$$

which is of the correct order of magnitude, but differs from the observed value of $6.674\,28(67) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$ [5, 6] by a substantial margin of 33σ .

C. Radiative corrections

The greatest constraint on the accuracy of the above calculation of G is not the accuracy to which the constituent electroweak parameters are known, but rather that the calculation was performed evaluating interaction (88) only to lowest order. One loop electromagnetic corrections arise from the diagrams of Fig. 11, and the resulting expression for G is

$$\begin{aligned} G &= \frac{e^4(1+a_e)}{8\varepsilon_0^2 m_e^4 T^2} \frac{\hbar}{c^5} \left[1 + \frac{3\alpha}{2\pi} + O(\alpha^2) \right] \\ &= 6.675\,514\,4(15) \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}, \end{aligned} \quad (116)$$

where α is the fine structure constant,

$$\alpha = \frac{e^2}{4\pi\varepsilon_0\hbar c}. \quad (117)$$




FIG. 11: One-loop diagrams making non-zero contributions to the value of G . Diagrams (i), (ii), and (iii) each acquire an additional factor of $\alpha/(2\pi)$ compared with Fig. 8, giving an $O(\alpha)$ correction factor of $[1 + 3\alpha/(2\pi)]$. Diagrams similar to (i) and (ii) but with the loop on the H or $H^\dagger$ boson legs do not yield any contribution.

Already our accuracy is much improved, with (116) differing from the experimental value by a mere 1.8σ (for $\sigma_{\text{NIST}} = 6.7 \times 10^{-15} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$). Again the leading constraint on our accuracy is lack of knowledge of higher order corrections due to diagrams involving two or more loops. However, the two-loop correction is anticipated to be of order $10^{-16} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$, and corrections beyond the two-loop level are probably unimportant given current precisions for e , $\hbar$, and m_e . The impact of these further corrections on this result will therefore be small.

Note also that the value of G computed is independent of our choice of convention for boson number density. A change

$$\int dt \|b_\mu\| = 1 \longrightarrow \int dt \|b_\mu\| = k \quad (118)$$

$$\Rightarrow g \longrightarrow \frac{g}{k} \quad (119)$$

will introduce a factor of k^{-2} from g^2 but a factor of k^2 from the fields corresponding to single H and $H^\dagger$ bosons in Fig. 8. Consequently the value of G remains

unchanged. The reader may wonder at the origin of an apparent new constant in the form of T^2 in the denominator of (115): This factor comes from g^2 , and its origin is simply in the definition of interaction strength from the boson-mediated coupling between two *single* electrons. It has no profound physical meaning, but is the part of g which reflects our definition of particle number density, scaling as k under transformation (118-119).

D. Prospects for detection of $G^{(\dagger)}$ bosons

For the spherically symmetric example considered, it was possible to eliminate all $G^{(\dagger)}$ interactions on macroscopic length scales, outside of a central singularity in the $G^{(\dagger)}$ field which became a singularity in the induced space-time. Can systems exist for which this is not possible, and in which interactions with $G^{(\dagger)}$ bosons may take place? A possible candidate for such a system would be a perfectly symmetric, rapidly spinning stellar collapse. Classically, such a system has been argued to exhibit a naked singularity: A singularity in space-time curvature without an event horizon, from which photons could apparently escape. In this theory the space-time singularity would correspond to a singularity in the induced co-ordinate frame such that interactions with the $G^{(\dagger)}$ fields cannot be completely eliminated over some region. Particles entering this region could then interact with the $G^{(\dagger)}$ field in a manner which would be detectable in the curved space-time.

Arguably no other, less pathological situations permitting the detection of $G^{(\dagger)}$ bosons exist: Such situations would entail obligate smooth continuation of the space-time metric in a manner not consistent with elimination of the foreground $G^{(\dagger)}$ fields. This would imply either a sharp discontinuity in the G and $G^\dagger$ fields, or an asymmetry between the two. The former is penalised by an infinitely large contribution to the Lagrangian, and so is unphysical, and the latter is prohibited by our final choice of gauge in Sec. II A. Specifically, any G or $G^\dagger$ boson will sustain a virtual $\bar{\phi}_1 \bar{\sigma}^\mu \phi_3$ or $\phi_3 \bar{\sigma}^\mu \phi_1$ field respectively. Even if these fields are not probed in a given Feynman diagram, any asymmetry is still prohibited by (12), and smooth continuation of this choice of gauge requires that the surplus particle is not a $G^{(\dagger)}$ boson but a $W^{(\dagger)}$ boson. Ultimately this gauge transformation rotates the $SU(2)$ subspace at the space-time co-ordinate where the isolated $G^{(\dagger)}$ boson would have been generated, mixing all interactions occurring at that point such that G and $G^{(\dagger)}$ bosons (if any) are produced in equal measure. Thus (12) prohibits production of isolated G or $G^\dagger$ bosons, as well as permitting elimination of the right-handed weak interaction.

In conclusion, detection of $G^{(\dagger)}$ bosons, except implicitly as the mediators of space-time curvature, is therefore impossible outside of a naked space-time singularity, and their interactions may be safely ignored.

E. Weak interactions involving $G^{(\dagger)}$ bosons

Interactions which ostensibly produce a single $G^{(\dagger)}$ boson may still take place, for example

$$W^+ + H^{--} \longrightarrow G^- + A, \quad (120)$$

but only if they are accompanied by a second reaction producing a G boson of the opposite type at the same space-time co-ordinate, e.g.

$$W^- + H^{++} \longrightarrow G^+ + A. \quad (121)$$

In fact, with g much reduced from the standard model, the paired interactions (120-121) are responsible for nearly all observed $W^+ + W^- \rightarrow 2A$ annihilation events. Examining the W and $W^\dagger$ bosons in (120-121), each participates in an interaction with cross-section $g^2/\sqrt{8}$, but the diagrams are augmented by an additional factor $\langle \|H_\mu^{(\dagger)}\| \rangle$ corresponding to the number of available H or $H^\dagger$ bosons in the background fields. Being unaware of the interactions between $H^{(\dagger)}$ and $G^{(\dagger)}$ bosons, we inadvertently conflate these processes with

$$W^+ + W^- \longrightarrow 2A, \quad (122)$$

giving an apparent W - $W^\dagger$ coupling

$$\frac{g_{\text{effective}}^2}{4} \equiv \frac{g^2 \langle \|H_\mu^{(\dagger)}\| \rangle T}{\sqrt{8}}. \quad (123)$$

The effective interaction strength is augmented by the number density of the background $H^{(\dagger)}$ fields, and by a factor of $\sqrt{2}$ due to the different boson-boson interactions involved. Together these effects offset the reduced value of g in the present model, and may be compared with the role of the background photon field in Sec. IV B.

A similar process enhances the strength of the $W^{(\dagger)}$ - Z coupling, but no equivalent exists for the $W^{(\dagger)}$ - N coupling, which is therefore negligible in the $r \gg r_{\text{crit}}$ regime.

VI. DEVIATION FROM GENERAL RELATIVITY

The above derivation of the Schwarzschild metric holds only in the combined limits

$$r \gg \frac{2GM}{c^2} \quad (124)$$

$$r \gg \sqrt{\frac{e^4 M \hbar}{16 \varepsilon_0^2 \pi m_e^4 T (1 + a_e)^2 c^6}}, \quad (125)$$

where the latter expression corresponds to r_{crit}^H for a body of mass M . The former corresponds to the Schwarzschild radius, indicating that this theory will predict novel physics for bodies classically presumed to collapse into black holes, and the latter corresponds to a radius below

which the role of the background field becomes less significant, the net generation of $G^{(\dagger)}$ pairs at the expense of the $H^{(\dagger)}$ fields decreases, and the effective value of G consequently becomes less.

Because these limits express different dependencies on M , we see that for

$$M > 1.5 \times 10^4 \text{ kg} \quad (126)$$

the physics of the theory begins to deviate from general relativity only on approaching the Schwarzschild radius. For suitably compressed bodies of mass $M < 1.5 \times 10^4 \text{ kg}$, however, the second limit is encountered first and there exists the tantalising prospect of observing a reduced gravitational interaction at radii less than r_{crit}^H . In practice this regime appears presently inaccessible, with r_{crit}^H being of order 10^{-23} m for the threshold mass, and scaling as $\sqrt{M}$.

VII. CONCLUSION

This paper has demonstrated the existence of a renormalisable extension of the standard model which is shown to reproduce some of the behaviours of the gravitational interaction for an irrotational, spherically symmetric point mass, most notably the Schwarzschild metric and the value of G , the Newtonian gravitational constant. That it is possible to construct such a theory suggests that there is profitable work on quantum gravity to be done in the low-energy limit, and that it may be possible to determine the form to which higher energy theories such as string theory must reduce.

The following outcomes appear to be relatively robust properties of this theory:

- The symmetry breaking process is more involved than the minimal scalar Higgs mechanism employed in the standard model.
- The symmetry group of the electroweak interaction must be expanded at least to $SU(3)$.
- The theory incorporates non-existence of neutrinos of right-handed helicity by construction.
- The neutrino is not its own anti-particle, implying non-occurrence of neutrinoless double beta decay.

That the theory is predictive of the value of G is already a significant test of the hypothesis that this is a valid representation of the gravitational interaction at low energies. Undoubtedly, as the theory is explored in more detail other opportunities to test this hypothesis will arise.

Finally, it is interesting to note that the theory includes a natural candidate for a dark matter particle, the N boson, which has a mass similar to that of the W and Z bosons. Unfortunately no prediction can yet be made about the interaction cross-section with nucleons, which is typically probed in dark matter detection experiments, as the theory presented here does not yet incorporate the strong nuclear force.

APPENDIX: STRUCTURE CONSTANTS OF $GL(3, \mathbb{R})$

When performing computations involving boson-boson interactions, it is useful to know the structure constants for a basis of $GL(3, \mathbb{R})$ which corresponds to the bosons being studied. Consider the basis

$$\begin{aligned}
 C_1 &= \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} & C_2 &= \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
 C_3 &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} & C_4 &= \frac{2}{\sqrt{6}} \begin{pmatrix} -\frac{1}{2} & 0 & 0 \\ 0 & -\frac{1}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
 C_5 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} & C_9 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
 C_6 &= C_2^\dagger & C_7 &= C_3^\dagger & C_8 &= C_5^\dagger,
 \end{aligned} \tag{127}$$

which satisfies $\text{Tr } C_a C_a^\dagger = 1$ (no sum over a). Identifying

$$\begin{aligned}
 b_\mu^1 &= A_\mu & b_\mu^2 &= H_\mu & b_\mu^3 &= G_\mu \\
 b_\mu^4 &= Z_\mu & b_\mu^5 &= W_\mu & b_\mu^6 &= N_\mu \\
 g &= \sqrt{2} f.
 \end{aligned} \tag{128}$$

the non-zero structure constants for this basis are

$$\begin{aligned}
 1/\sqrt{2} &: k_{133}, k_{188}, k_{371}, k_{515}, k_{717}, k_{851} \\
 -1/\sqrt{2} &: k_{155}, k_{177}, k_{313}, k_{581}, k_{731}, k_{818} \\
 1 &: k_{253}, k_{382}, k_{393}, k_{576}, k_{595}, k_{635}, \\
 &\quad k_{728}, k_{867}, k_{977}, k_{988} \\
 -1 &: k_{278}, k_{365}, k_{523}, k_{687}, k_{756}, k_{797}, \\
 &\quad k_{832}, k_{898}, k_{933}, k_{955} \\
 3/\sqrt{6} &: k_{343}, k_{477}, k_{488}, k_{545}, k_{734}, k_{854} \\
 -3/\sqrt{6} &: k_{374}, k_{433}, k_{455}, k_{584}, k_{747}, k_{848} \\
 \sqrt{2} &: k_{122}, k_{261}, k_{616} \\
 -\sqrt{2} &: k_{166}, k_{212}, k_{621}
 \end{aligned} \tag{129}$$

where

$$\sum_c k_{abc} C_c = [C_a, C_b]. \tag{130}$$

-
- [1] J. Polchinski, *String Theory*, vol. 1 & 2 (University Press, Cambridge, 2005).
 - [2] M. B. Green, J. H. Schwarz, and E. Witten, *Superstring Theory*, vol. 1 & 2 (University Press, Cambridge, 1988).
 - [3] F. Cianfrani, O. Lecian, and G. Montani, arXiv:0805.2503v2 [gr-qc] (2008).
 - [4] C. Rovelli, *Quantum Gravity* (University Press, Cambridge, 2007).
 - [5] P. J. Mohr, B. N. Taylor, and D. B. Newell, Rev. Mod. Phys. **80**, 633 (2008).
 - [6] P. J. Mohr, B. N. Taylor, and D. B. Newell, J. Phys. Chem. Reference Data **37**, 1187 (2008).
 - [7] L. H. Ryder, *Quantum Field Theory* (University Press, Cambridge, 1996), 2nd ed.
 - [8] S. Weinberg, Phys. Rev. Lett. **19**, 1264 (1967).
 - [9] J. Wess and J. Bagger, in *Gauge Theories in High Energy Physics*, edited by M. K. Gaillard and R. Stora (North Holland, Amsterdam, 1983), vol. 37, pp. 1–57.
 - [10] C. Amsler, M. Doser, M. Antonelli, D. M. Asner, K. S. Babu, H. Baer, H. R. Band, R. M. Barnett, E. Bergren, J. Beringer, et al., Physics Letters B **667**, 1 (2008).
 - [11] A. Goobar, S. Hannestad, E. Mortsell, and H. Tu, Journal of Cosmology and Astroparticle Physics **2006**, 019 (2006).
 - [12] S. Coleman and J. Mandula, Phys. Rev. **159**, 1251 (1967).